



# MULTI-SPECIES HOUSING PROTOCOL

(i.e. Husbandry & Education Buildings)

## Footbaths:

1. Fill with a 1-ounce per gallon solution of Animal Facility Disinfectant.
2. Refresh footbath solution every 24-hours, or when the solution becomes visibly dirty.
3. Footbath solution should be changed multiple times a day in isolation areas.

## Mews & Chickens:

1. Use STABLE Environment ONLY!
2. Rake and sweep to remove all loose bedding and debris.
3. Pour full strength **STABLE Environment** into Foam Gun reservoir tank.
4. Refer to Foam Gun Tip Chart to select dilution. *Recommended dilution: 2 or 4-ounces per gallon of water.*
5. Attach reservoir tank to spray fitting.
6. For stones and substrate apply every other day. Allow the solution to stand for 3 to 5 minutes and rinse.
7. For perches and nesting boxes, dilute STABLE Environment in a bucket and apply using a brush. Allow the solution to stand for 3 to 5 minutes and rinse.
8. Allow solution to stand for 3 to 5 minutes.
9. **Monthly Deep Clean:** Remove birds from the area and use foam gun to apply STABLE Environment to all surfaces.
10. Agitate surface if needed.
11. Rinse the perches and nesting boxes before putting birds back into the area.
12. Sweep or squeegee surface to remove any puddles before returning birds back to the area.

## Bird & Small Animal Cages (excluding amphibians):

To minimize the migration of chemical sprays and vapors, when cleaning bird cage surfaces, we recommend using **wet wipes** in place of traditional spray bottles. If spray bottles are used, set the sprayer setting to course/stream.

1. Prepare the **Refillable Wiping System**, by mixing 1-ounce of **Animal Facility Disinfectant** and 32-ounces of water.
2. Remove bird(s), food and water dishes from cage.
3. Remove and discard solid waste, bedding and other debris.
4. Wipe down surfaces including perches, play structures, etc..
5. Scrub surface with a **synthetic brush**.
6. Allow solution to stand for 10-minutes.
7. Use a wet sponge or cloth to remove any heavy moisture and emulsified soil.
8. Allow surfaces to air dry before returning animals to the cage.
9. **NOTE:** For large, multi-bird enclosures, use a **foam gun applicator** or pump sprayer to apply the solution. For heavily soiled surfaces prepare and use a solution of **Pro Foam Ultra** to preclean the surface before implementing the disinfecting process.



# MULTI-SPECIES HOUSING PROTOCOL

(i.e. Husbandry & Education Buildings)

## Show & Daily Use Crates:

1. Spot clean using a 1-ounce per gallon solution of Animal Facility Disinfectant and water.
2. Twice weekly use Animal Facility Disinfectant in a foam gun and apply to interior and exterior surfaces and crate mats. Allow the solution to stand for 10-minutes and rinse thoroughly.
3. Allow surfaces to dry thoroughly before using.

## Program & Off-Site Crates:

1. Rinse to remove any fecal matter or other soil from both the interior and exterior surface.
2. Use an Animal Facility Disinfectant Refillable Wipe to wipe down the interior and exterior surface.
3. For crates that are too deep to reach by hand, use a bucket and brush to scrub the surfaces.
4. Allow surfaces to dry thoroughly before use.

## Amphibian & Arthropod Carriers:

1. DO NOT USE Animal Facility Disinfectant!
2. Use a 10% bleach solution to clean the carrier.