

ProVetLogic

PROTOCOL



ProVetLogic Professional & ProLoc Wall-Mounted Dispenser

Product Dilution & Use Efficacy

GREEN 1.25 OZ PER GALLON OF WATER

CANINE

Kennel Cough
Canine Adenovirus
Canine Coronavirus
Canine Distemper
Canine Parainfluenza
Canine Hepatitis
Infectious Bronchitis
Rabies Virus

FELINE

Feline Calicivirus
Feline Peritonitis
Feline Leukemia
Feline Panleukopenia
Feline Picornavirus
Feline Rhinotracheitis

AVIAN

Avian Influenza A H9N2
Avian Laryngotracheitis
Avian Influenza A H5N1
Avian Reovirus

FARM

Equine Arteritis Virus
Influenza A2/Japan
Newcastle Disease
Porcine Rotavirus
Pseudorabies Virus
Porcine Respiratory
Infectious Bovine
Transmissible (TGE)

OTHER

Aspergillus Niger
Trichophyton Menta
MRSA
E-Coli and Salmonella
H1N1

RED 4 OZ PER GAL

Canine Parvovirus
Minute Virus of Mice
Porcine Parvovirus

RECOMMENDED TOOLS FOR THE JOB



Labeled Bottles



Pump-Sprayer or Foam Gun



Mop Bucket



Synthetic Tools



Safety Signs

HOSPITAL HARD SURFACE & FLOOR CLEANING/DISINFECTING



1

Remove all animals from the immediate area. Return after all visible puddles and heavy moisture are removed from surface.



4

Apply solution liberally over the entire surface. Allow to dwell for at least 10 minutes, agitate if necessary and wipe away access moisture. Allow to air dry.



2

Remove all bedding, shavings, toys, food and water dishes, as well as feces and urine.



5

Fill mop bucket with a prepared solution of product and water. **DO NOT FILL TO THE TOP!** Empty when water becomes visibly dirty and rinse bucket and mop.



3

Mix solution in a labeled spray bottle and apply using coarse spray. Mist setting may cause URI in cats and birds.



6

Start from the back of the room, working to the front. Remove any excess puddles and moisture and allow surface to air dry.

KENNEL & LARGE ANIMAL CONTAINMENT CLEANING/DISINFECTING



1

Remove all animals from the immediate area. Return after all visible puddles are removed.



4

Use a Pump-Up Sprayer or Automatic Spray/Foam Gun to apply the solution. Start from the top of area, working down towards the bottom and from back to front.



2

Remove all bedding, shavings, toys, food and water dishes, as well as feces.



5

Using a brush and handle made of synthetic materials, agitate the soiled surfaces, including walls, floors, cage doors and hinges. Allow solution to dwell at least 10 minutes.



3

Rinse surface to remove any loose debris, feces, etc. Avoid heavy pressure, as this may spread infectious agents.



6

Use clean water to rinse away any remaining soap and soil residue and allow the surface to air dry. Return animals only after all visible puddles are removed.

ProVetLogic Professional – 800.869.4789 or Visit Us at www.provetlogic.com